

Semi-Annual

REPORT ON J-209

GRANT
CONTRACT NO DA -CRD-AG-S92-544-64-G23

INCLUSIVE DATES 29 July 1964 TO 28 Jan. 1965

AD625955

SUBJECT OF INVESTIGATION

HISTOCHEMICAL STUDIES ON THE DISTRI-
BUTION OF ENZYMES, ESPECIALLY OXIDASES
AND PHOSPHATASES IN THE LIVING BODY

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION			
Microcopy	Microfilm		
1.00	0.50	7	as
AR 10000 COPY			


Call 1

RESPONSIBLE INVESTIGATOR

Tadao Mitsui, M.D.

Professor of Anatomy
School of Medicine, Keio University
Shinjuku-ku, Tokyo, Japan

PROCESSED COPY



U.S. Army Research & Development Group (998) (Far East)

Office of the Chief of Research and Development

United States Army

~~APG-543~~
APO San Francisco 96343

Best Available Copy

D-I-S-T-R-I-B-U-T-I-O-N

The distribution of this report as made by USA R&D Op (FE)
is as follows:

Assistant Director of Army Research (4)
Office, Chief of Research and Development, DA
ATTN: Chief, Research Programs Office
3045 Columbia Pike, Arlington, Virginia 22204

Army Attache, American Embassy, Tokyo, Japan (1)

~~U. S. Army Medical Research and Development Command, (-)~~
~~Washington, D. C. 20315~~

~~Defense Documentation Center, Cameron Station, (-)~~
~~Alexandria, Virginia. ATTN: TISIA-2~~

Office of Primary Scientific Liaison (4)
U.S. Army Medical Research and Development Comd
ATTN: Chief, Medical Research Branch
Washington, D. C. 20315

Office of Scientific Cognizance (2)
Scientific Director
Armed Forces Institute of Pathology
Washington, D. C.

Best Available Copy

Semifinal Report

DA-CRD-AG-S92-544-64-G23

29 July 1964 to

28 Jan. 1965

Histochemical studies on the distribution of
enzymes, especially oxidases and phosphatases
in the living body

Tadao Mitsui, M.D.

Professor of Anatomy, School of Medicine
Keio University, Shinjuku-ku, Tokyo
Japan

Histochemical studies on the distribution
of enzymes, especially oxidases and
phosphatases in the living body

Tadao Mitsui, M.D.

Professor of Anatomy, School of
Medicine, Keio University
Shinjuku-ku, Tokyo, Japan

A B S T R A C T

Two giant salamanders, *Megalobatrachus japonicus* were used for materials.

The blood was carefully taken from the tail and smear preparations were obtained. The blood smears were fixed one minute with ethanol-formol (9:1) solution prior to staining, washed thoroughly in distilled water, and stained five minutes with either of the following peroxidase reagents freshly prepared each time.

1) Mixture of benzidine 0.3 gr., 3% H_2O_2 10 drops, and distilled water 100 cc. This mixture should be filtered because of poor solubility of benzidine in water.

2) Mixture of orthophenylenediamine 0.1 gr., 3% H_2O_2 10 drops, and distilled water 100 cc.

The smears were then washed in water, and counterstained with dilute G i e m s a stain for ten to fifteen minutes.

Finally they were washed in water, and exposed to air till dry.

The alkaline phosphatase technic (modified technic of the leadsulphide method of T a k e u c h i) was also applied for blood staining.

The preparation stained with the benzidine peroxidase reagent, that with the orthophenylenediamine reagent, and that with G i e m s a stain, were simultaneously observed, and compared.

In the giant salamander, the neutrophil leukocytes generally showed peroxidase activity, while the eosinophil and basophil leukocytes did not. The reaction was lacking in all cells of the lymphatic series. As is well known, the general feature of the peroxidase reaction in human leukocytes under the light microscope is that the cytoplasm is diffusely, fine granularly stained except eosinophil leukocytes in which the reaction may take a pattern of prominent coarse granules corresponding to the eosinophil granules.

A gigantic peroxidase granule was detected in the polymorphonuclear neutrophil leukocytes of the giant salamander, *Megalobatrachus japonicus*. Among 70 different animals examined, this gigantic peroxidase granule was found only in the giant salamander in apparently healthy condition. This granule was generally localized in the center of the leukocyte, single in number, six to ten μ in diameter, and could be demonstrated both with the benzidine peroxidase

reagent and the orthophenylenediamine peroxidase reagent.

This feature resembles the gigantism of the peroxidase granule that was recognized in the C h e d i a k (1952) and H i g a s h i 's disease (1954). This congenital human disease is characterized by albinism with photophobia, early death of albinic sibs and peculiar pathological blood picture in which the above-mentioned gigantic granules or inclusion bodies are demonstrated in the leukocytes. In this disease the peroxidase granules gather in a few or several spots to form round clumps or globes not only in the neutrophils but also in the monocytes, eosinophils, myelocytes and the myeloblasts, in other words it is an abnormality affecting all the cells of the myelogenous system.

It is of interest that one of pathological characters of human leukocytes was found in an apparently healthy giant salamander, although it remains to be determined in what way this gigantic peroxidase granule developed only in the neutrophil leukocytes of this animal.

The gigantic peroxidase granule in the giant salamander may be derived from a conglomeration of the peroxidase substance within the leukocyte under abnormal or normal metabolism of protein and other substances in

this animal, or may be a product caused by phagocytosis of a peroxidase reacting substance. However, if a phagocytosis phenomenon is assumed in the giant salamander, it remains still unanswered why it occurred only in the neutrophil leukocytes and not at all in other leukocytes such as monocytes.

It is of special interest that peculiar and significant findings in blood cells of man and animals may frequently be obtained with the aid of these cytochemical reactions.